



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Foxes As a Potential Wildlife Reservoir for mecA-Positive Staphylococci

**Citation for published version:**

Carson, M, Meredith, AL, Shaw, DJ, Giotis, ES, Lloyd, DH & Loeffler, A 2012, 'Foxes As a Potential Wildlife Reservoir for mecA-Positive Staphylococci' *Vector Borne and Zoonotic Diseases*, vol. 12, no. 7, pp. 583-587. DOI: 10.1089/vbz.2011.0825

**Digital Object Identifier (DOI):**

[10.1089/vbz.2011.0825](https://doi.org/10.1089/vbz.2011.0825)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

*Vector Borne and Zoonotic Diseases*

**Publisher Rights Statement:**

This is a copy of an article published in the *Vector-Borne and Zoonotic Diseases* ©2012 [copyright Mary Ann Liebert, Inc. *Vector-Borne and Zoonotic Diseases* is available online at: <http://online.liebertpub.com>.

<http://www.liebertpub.com/nv/resources-tools/self-archiving-policy/51/>

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## Foxes As a Potential Wildlife Reservoir for *mecA*-Positive Staphylococci

Marianne Carson,<sup>1</sup> Anna L. Meredith,<sup>2</sup> Darren J. Shaw,<sup>2</sup> Efstathios S. Giotis,<sup>1</sup>  
David H. Lloyd,<sup>1</sup> and Anette Loeffler<sup>1</sup>

### Abstract

Methicillin-resistant staphylococci (MRS), and methicillin-resistant *Staphylococcus aureus* (MRSA) in particular, have become a public and veterinary health concern. The search for MRS reservoirs outside human hospitals is needed in order to understand the reasons for their persistence and to control their spread. MRS have been isolated from rats, but little is known about their occurrence in foxes. In view of the perceived increasing proximity between people and foxes in the U.K. and the well-documented potential of foxes as hosts for zoonotic pathogens, this study examined whether foxes can be a reservoir for MRS. This study examined the carriage of staphylococci and their antimicrobial resistance patterns in 38 foxes (*Vulpes vulpes*) from rural and semirural areas in the U.K. Staphylococci were isolated by enrichment culture from nasal, oral, axillary, and perineal swabs and speciated by standard bacteriological tests and API ID32 STAPH (bioMérieux, Marcy l'Étoile, France). Antimicrobial resistance was investigated by disc diffusion tests and identification of *mecA*. Thirty-seven staphylococcal isolates were identified from 35 of the 38 foxes. All isolates were coagulase-negative and most frequently included species from the *S. sciuri* group (35%), *S. equorum* (27%), and *S. capitis* (22%). All were phenotypically resistant to methicillin, and *mecA* was detected in 33 (89%) of isolates, but only 10 (27%) showed broad  $\beta$ -lactam antibiotic resistance. Methicillin-resistant *S. aureus* was not identified. These results indicate that foxes are a potential wildlife reservoir for *mecA*-positive staphylococci. Selection pressure from environmental contamination with antimicrobials should be considered.

**Key Words:** Antimicrobial resistance—Foxes—*mecA*—Staphylococci—Wildlife.

### Introduction

ANTIMICROBIALS ARE WIDELY USED FOR THERAPY and prevention of disease in human and veterinary medicine as well as in modern agriculture (Cabello 2006; Owens and Stoessel 2008). Expansive use has led to selection for antimicrobial resistance in bacteria, the spread of associated resistance genes, and to the emergence of multidrug-resistant pathogens that pose a potential threat to human health (Baquero et al. 2009; Tacconelli 2009; Heuer et al. 2011). Staphylococci, which colonize and infect humans and animals, and which can persist on environmental surfaces for several months, have been known for their potential to express antibiotic resistance since 1944 (Kirby 1944). Furthermore, they have been associated with multidrug-resistance since the emergence of methicillin-resistant *Staphylococcus aureus* (MRSA) in human hospitals. Methicillin resistance has also

been reported in up to 80% of clinical isolates of coagulase-negative staphylococci (CNS), and seems to be associated with concurrent resistance to non- $\beta$ -lactam antimicrobial classes such as to aminoglycosides, macrolides, quinolones, and tetracyclines (Hope et al. 2008). Important examples of CNS in human medicine include *S. epidermidis*, *S. lugdunensis*, and *S. saprophyticus*, which have emerged as the most common nosocomial pathogens isolated from bloodstream infections in several countries, and are frequently associated with implant- or catheter-related infections in immunocompromised people, or with endocarditis and urinary tract infections in immunocompetent hosts (Piette and Verschraegen 2008).

Resistance to methicillin is conferred by the *mecA* gene, which is integrated into the bacterial chromosome on a large mobile cassette element (SCC*mec*), and confers broad  $\beta$ -lactam antibiotic resistance via an altered penicillin-binding protein

<sup>1</sup>Department of Veterinary Clinical Sciences, Royal Veterinary College, University of London, London, United Kingdom.

<sup>2</sup>Royal (Dick) School of Veterinary Studies, University of Edinburgh, Edinburgh, United Kingdom.

(De Lencastre et al. 2007). *mecA* in MRSA is thought to have been acquired by *S. aureus* through horizontal transfer from coagulase-negative species, and a wildlife origin for *mecA* has been proposed. Genetic homologues of the early *S. aureus* SCC*mec* elements have been identified in *S. sciuri*, and biochemical similarities between their penicillin-binding proteins have been demonstrated (Couto et al. 1996; Chambers 1997; Wu et al. 2001). The results from genetic analyses have since led to taxonomic changes, and due to the close relatedness of the staphylococcal species *S. sciuri*, *S. vituli*, and *S. lentus*, these three are now combined in the *S. sciuri* group. Members of this group are frequently found in rodents and other mammals, but also in humans (Stepanovic et al. 2005).

The search for MRS reservoirs outside human hospitals continues in order to understand and predict the epidemiology of antibiotic resistance, and to help limit the spread of methicillin-resistant pathogens. Companion animals and more recently pigs have been linked to recurrent MRSA outbreaks in human hospitals, and genetic analyses have implicated in-contact animals as vectors for human MRS acquisition (Voss et al. 2005; Loeffler and Lloyd 2010). Similarly, MRS have been identified from indoor and outdoor environmental sources (Scott et al. 2008; Soge et al. 2009).

However, little has been reported on the occurrence of MRS in wild animals, despite several reports highlighting examples of wildlife as reservoirs for antibiotic resistance. Antimicrobial resistance in staphylococci was higher in isolates from feral cats in the U.K. than from domestic cats (Patel et al. 1999). Rats on swine farms carried MRSA ST398, and were considered potential contributors to the spread of resistance within and between swine herds (van de Giessen et al. 2009). Multiple antimicrobial resistances were found in *Escherichia coli* isolated from small wild animals living in proximity to swine farms, and extended-spectrum  $\beta$ -lactamase-producing *E. coli* was isolated from foxes in central Europe (Literak et al. 1984; Kozak et al. 2009). Wild red-billed choughs (Family *Corvidae*) living in areas where manure land spreading is practiced showed high levels of multidrug resistance in their bacterial flora, in accord with resistance levels found in the manure (Blanco et al. 2009). Wild-living primates in contact with human refuse showed higher levels of resistant microflora compared to primates with little or no contact with humans (Rolland et al. 1985).

Following the dramatic expansion of red fox populations in some countries in Europe over the past decade, and the perceived closer proximity of foxes to humans (Gloor et al. 2001; Canid Specialist Group 2004), this study aimed to determine whether foxes should be considered a potential wildlife reservoir or vector for MRSA and other MRS.

## Materials and Methods

### Sampling of foxes

Thirty-eight dead foxes were sampled for staphylococcal carriage in the Scottish Borders and Pentlands between May and November 2007 and June and September 2008. The foxes were shot as part of pest-control programs in these areas, but the carcasses were retained for a Department for Environment, Food and Rural Affairs (DEFRA)-funded study into the role of carnivores as sentinels for emerging diseases being carried out at the Royal (Dick) School of Veterinary Studies. The animals were from rural areas except for four, which were found in semi-rural agricultural regions. They included male

and female adult and subadult foxes, with some juveniles. The animals were sampled within 72 h of death.

Four sites were sampled per animal, including the nostrils (both with the same swab), buccal mucosa, axillary skin, and the perianal skin. The swabs were rolled over skin or mucosae for approximately 5 sec, and posted to the Royal Veterinary College in charcoal transport medium.

### Swab processing

The four swabs from each animal were aseptically pooled in 10 mL tryptone soya broth (Oxoid, Basingstoke, Hampshire, U.K.) supplemented with sodium chloride (Sigma-Aldrich, Gillingham, Dorset, U.K.), to a total salt concentration of 10% for selective enrichment of staphylococci. After 24 h incubation at 37°C, the samples were streaked out onto 5% sheep blood agar (Oxoid), and presumptive staphylococci were identified based on salt resistance, colony morphology, Gram staining characteristics, and a positive catalase test. Up to three morphologically distinct staphylococcal colonies per sample were frozen at -80°C in brain heart infusion (Oxoid) with 20% glycerol (Sigma-Aldrich).

### Staphylococcus species identification

Staphylococci were regrown from frozen and incubated at 37°C on blood agar for 24 h, and on mannitol salt agar (both Oxoid) and mannitol salt agar with 6 mg/L oxacillin (Sigma-Aldrich) for 48 h. Phenotypic tests for speciation included assessment of hemolysis, a slide coagulase test with dog plasma, and where negative, a tube coagulase test with rabbit plasma, DNase test, and acetoin fermentation (Voges-Proskauer reaction; Barrow and Feltham 1993). The API ID32 STAPH (bioMérieux, Marcy l'Etoile, France) identification kit was used for further speciation following the manufacturer's instructions. API strips were read manually and species identification was based on interpretation of scores by bioMérieux's online *apiweb*<sup>TM</sup> software.

### Phenotypic resistance profiles

Methicillin resistance was screened for by growth on oxacillin-supplemented mannitol salt agar as described above, and agar color change from pink to yellow was indicative of mannitol fermentation. Antimicrobial resistance profiles were determined by disc diffusion tests on Mueller-Hinton agar following the Clinical and Laboratory Standards Institute protocol (The Clinical and Laboratory Standards Institute 2004) with colonies grown on blood agar. The following antimicrobial discs were used (Oxoid): methicillin (5  $\mu$ g), ampicillin (10  $\mu$ g), amoxicillin/clavulanic acid (30  $\mu$ g), cefalexin (30  $\mu$ g), clindamycin (2  $\mu$ g), tetracycline or oxytetracycline (both 30  $\mu$ g), trimethoprim-sulfamethoxazole (25  $\mu$ g), enrofloxacin (5  $\mu$ g), and fusidic acid (5  $\mu$ g). Breakpoints were as defined by CLSI where available, for fusidic acid as defined by Olsson-Liljequist and associates (2002), and for cefalexin and enrofloxacin at 12 mm, in accordance with the manufacturers' recommendations. Multidrug-resistance was defined as resistance to three or more non- $\beta$ -lactam antimicrobial classes (Merlino et al. 2002).

### Genotypic resistance to methicillin

The presence of *mecA* was investigated in all isolates that were phenotypically resistant to methicillin by gene

amplification through polymerase chain reaction (PCR) and subsequent gel electrophoresis. DNA was extracted from overnight cultures using a commercial purification kit (Bacterial Genomic DNA Purification kit; Edge Biosystems, Gaithersburg, MD) according to the manufacturer's instructions. Samples were prepared for PCR by adding 1  $\mu$ L of extracted DNA to 49  $\mu$ L PCR mix. The mix contained 5  $\mu$ L 10 $\times$ PCR buffer, 1  $\mu$ L 10 mM dNTP mix, 2  $\mu$ L each of forward and reverse *mecA* primers, and 0.25  $\mu$ L HotStarTaq DNA Polymerase (Qiagen, Crawley, West Sussex, U.K.). PCR primers and conditions were as described by Brakstad and colleagues (1992). A *S. aureus* MRSA ST398 t011 was used as *mecA*-positive control for all phenotypic tests and the PCR.

## Results

### Staphylococcus isolation and species identification

Thirty-five of the 38 foxes (92.1%) yielded staphylococci, and 37 different isolates were investigated. Two morphologically distinct staphylococcal isolates were identified from two foxes, and in both cases the two isolates represented two different species with distinct antimicrobial resistance patterns.

All isolates were coagulase-negative on both slide and tube coagulase testing. Two isolates were DNase-positive. These were also two of three isolates exhibiting  $\beta$ -hemolysis. Seven distinct staphylococcal species were identified by API ID32 STAPH. *S. sciuri* (35%), *S. equorum* (27%), and *S. capitis* (22%), were the most frequently identified species. Two isolates were identified as *S. chromogenes* (5%), two as *S. xylosus* (5%), one as *S. lentus* (3%), and another as *S. kloosi* (3%). All isolates were Voges-Proskauer negative, and these results matched the acetoin fermentation on API in all.

### Antimicrobial resistance

All isolates grew on oxacillin-screening agar, and all showed resistance to methicillin on disc diffusion tests. High levels of resistance were seen, particularly to the other  $\beta$ -lactam antibiotics (Table 1), but only 10 isolates showed resistance to all four  $\beta$ -lactam antibiotics (Table 2). For non- $\beta$ -

TABLE 1. FREQUENCY OF RESISTANCE TO DIFFERENT ANTIMICROBIAL AGENTS DETERMINED BY DISC DIFFUSION TESTING AMONG 37 STAPHYLOCOCCAL ISOLATES FROM FOXES

Class of antimicrobial	Antimicrobial disc	Resistant isolates (%)
$\beta$ -Lactam		
	Penicillin	
	Methicillin 5 $\mu$ g	37 (100)
	Ampicillin 10 $\mu$ g	25 (67.6)
	Amoxicillin/clavulanic acid 30 $\mu$ g	19 (51.4)
Cephalosporin	Cefalexin 30 $\mu$ g	17 (45.9)
Fluoroquinolone	Enrofloxacin 5 $\mu$ g	2 (5.4)
Potentiated sulfonamide	Trimethoprim-sulfamethoxazole 25 $\mu$ g	0 (0)
Tetracycline	Oxytetracycline and tetracycline 30 $\mu$ g	5 (13.5)
Lincosamide	Clindamycin 2 $\mu$ g	2 (5.4)
Sterol	Fusidic acid 5 $\mu$ g	7 (18.9)

TABLE 2. PHENOTYPIC ANTIMICROBIAL RESISTANCE PATTERNS AMONG 37 STAPHYLOCOCCAL ISOLATES FROM FOXES

Resistance profiles	Frequency (%)
MET	3 (8.1)
MET, CL	6 (16.2)
MET, AMP	3 (8.1)
MET, TET	3 (8.1)
MET, CL, AMP	1 (2.7)
MET, AMC, AMP	4 (10.8)
MET, AMP, ENR	1 (2.7)
MET, AMP, DA	1 (2.7)
MET, CL, AMC, AMP	6 (16.2)
MET, AMC, AMP, TET	2 (5.4)
MET, AMC, AMP, FD	3 (8.1)
MET, CL, AMC, AMP, FD	2 (5.4)
MET, CL, AMC, AMP, ENR, FD	1 (2.7)
MET, CL, AMC, AMP, DA, FD	1 (2.7)

MET, methicillin; CL, cefalexin; AMP, ampicillin; AMC, amoxicillin/clavulanic acid; TET, tetracycline; ENR, enrofloxacin; DA, clindamycin; FD, fusidic acid.

lactam antibiotics, fusidic acid (19%) resistance was most frequently seen, followed by tetracycline (14%). Low frequencies of resistance (5%) were seen for enrofloxacin and clindamycin. None of the isolates were resistant to trimethoprim-sulfamethoxazole. Fourteen distinct resistance patterns were observed, and 84% of isolates were resistant to at least two of the four  $\beta$ -lactams, and 54% were resistant to at least three (Table 2). Most isolates (62%) were susceptible to all non- $\beta$ -lactam antibiotics. In addition to  $\beta$ -lactam resistance, 32.4% showed resistance to one other class of antimicrobial, two isolates (5.4%) were resistant to two non- $\beta$ -lactam antibiotics, while multidrug-resistance was not observed.

### Presence of *mecA*

The presence of *mecA* was confirmed in 89% (n=33) of phenotypically methicillin-resistant isolates. The four *mecA*-negative isolates did not appear to be associated with a specific species or disc resistance pattern. Three of those were resistant to all four  $\beta$ -lactam antibiotics including cefalexin, and the fourth showed resistance to three  $\beta$ -lactam antibiotics and tetracycline, but not cefalexin. Two of the *mecA*-negative isolates were DNase-positive, showed  $\beta$ -hemolysis, and were identified as *S. chromogenes* by API.

## Discussion

The results from this study indicate that the staphylococcal microflora of foxes are a potential wildlife reservoir for *mecA*, a resistance gene of clinical significance in human medicine worldwide. Although a larger sample size and a more diverse geographical sampling area would have allowed statistical comparison between locations and proximity to people, to the authors' knowledge, this is currently the largest reported study on staphylococci and antimicrobial resistance in foxes. Foxes are well known vectors for zoonotic pathogens such as scabies mites and dermatophytes, but they have also been shown to carry mycobacteria (Millán et al. 2008), which may be associated with zoonotic spread and multidrug-resistance.

While all were negative for MRSA, the majority of foxes carried coagulase-negative MRS. High frequencies of coagulase-negative MRS have previously been reported in feral cats, and resistance had been attributed to environmental sources (Patel et al. 1999; Hariharan et al. 2011). In companion animals and people, coagulase-negative MRS had been isolated from between 20% and 82% of horses (Baptiste et al. 2005; Bagcigil et al. 2007; Loeffler et al. 2010), from 63% of their in-contact humans, and from 66% of stable environmental sites (Moodley and Guardabassi 2009). In contrast, only 7–15% of domestic dogs and cats were positive for coagulase-negative MRS in these screening studies. This may indicate that some hosts such as horses, humans, and now also foxes, support carriage of coagulase-negative MRS better than others, regardless of antimicrobial selective pressure.

All isolates in this study showed phenotypic resistance to methicillin, but broad-spectrum  $\beta$ -lactam resistance and *mecA* detection were not consistent in all. While procedural error cannot be excluded, similar findings have been reported in a CNS isolate (van Duijkeren et al. 2004), and most recently in the newly identified *mecA* (LGA151)-MRSA isolated from cattle in the U.K. and Denmark (García-Álvarez et al. 2011). Vigilance and a continuing effort to improve laboratory methods are warranted in order to avoid misidentification of important pathogens. The use of more discriminatory molecular methods for such isolates could improve confidence with regard to speciation, as even coagulase test results may be unreliable, as has been described for *S. (pseud)intermedius* (Cox et al. 1985; Zadoks and Watts 2009).

The CNS species identified from the foxes in this study and the predominance of the *S. sciuri* group reflect those commonly found in a wide variety of domestic and wild animals (Kloos et al. 1976; Kawano et al. 1996; Nagase et al. 2002; Devriese et al. 2009; Zhang et al. 2009). *S. sciuri*, *S. equorum*, *S. xylosum*, and *S. lentus* have all been isolated from the intestines of wild small animals such as shrews and voles (Hauschild 2001). *S. chromogenes* is associated with udder health problems in dairy cattle (De Vlieghe et al. 2003). *S. capitis* is a common commensal of mainly human skin which has recently gained importance as an opportunistic pathogen, and *S. kloosii*, closely related to *S. sciuri*, is found mainly in wildlife (Schleifer et al. 1984; Takahashi et al. 1999). Members of the *S. sciuri* group are rarely found in humans and seldom cause disease, indicating that host-pathogen interactions and host specificity play a role in staphylococcal carriage (Irlinger 2008).

In conclusion, foxes appear to be competent carriers of several multi-host staphylococcal species and a potential source of the *mecA* gene. While the selection pressure on the microflora of foxes is likely to reflect a combination of naturally occurring and anthropogenic antimicrobial agents (Martinez 2008), a better understanding of their origin is needed to eventually limit the spread of resistance genes into pathogenic bacteria.

### Acknowledgments

Anette Loeffler and Anna Meredith were supported by research grants from DEFRA (U.K.) for research into MRSA in animals, and for evaluating predators as sentinels for emerging diseases (DEFRA projects OD2019 and SE01526, respectively).

### Author Disclosure Statement

No competing financial interests exist.

### References

- Bagcigil FA, Moodley A, Baptiste KE, et al. Occurrence, species distribution, antimicrobial resistance and clonality of methicillin- and erythromycin-resistant staphylococci in the nasal cavity of domestic animals. *Vet Microbiol* 2007; 121: 307–315.
- Baptiste KE, Williams K, Willams NJ, et al. Methicillin-resistant staphylococci in companion animals. *Emerg Infect Dis* 2005; 11:1942–1944.
- Barrow GI, Feltham RKA. Characters of Gram-positive bacteria. In: *Cowan and Steel's Manual for the Identification of Medical Bacteria*, 3rd ed. Barrow GI, Feltham RKA, eds. Cambridge: Cambridge University Press, 1993:50–93.
- Baquero F, Alvarez-Ortega C, Martinez JL. Ecology and evolution of antibiotic resistance. *Environ Microbiol Rep* 2009; 1:469–476.
- Blanco G, Lemus JA, Grande J. Microbial pollution in wildlife: Linking agricultural manuring and bacterial antibiotic resistance in red-billed choughs. *Environ Res* 2009; 109:405–412.
- Brakstad OG, Aasbakk K, Maeland JA. Detection of *Staphylococcus aureus* by polymerase chain reaction amplification of the *nuc* gene. *J Clin Microbiol* 1992; 30:1654–1660.
- Cabello FC. Heavy use of prophylactic antibiotics in aquaculture: a growing problem for human and animal health and for the environment. *Environ Microbiol* 2006; 8:1137–1144.
- Canid Specialist Group. 2004. [http://www.canids.org/species/Red\\_fox.pdf](http://www.canids.org/species/Red_fox.pdf)
- Chambers HF. Methicillin resistance in staphylococci: molecular and biochemical basis and clinical implications. *Clin Microbiol Rev* 1997; 10:781–791.
- Couto I, de Lencastre H, Severina E, et al. Ubiquitous presence of a *mecA* homologue in natural isolates of *Staphylococcus sciuri*. *Microb Drug Resist* 1996; 2:377–391.
- Cox HU, Newman SS, Roy AF, et al. Comparison of coagulase test methods for identification of *Staphylococcus intermedius* from dogs. *Am J Vet Res* 1985; 46:1522–1525.
- De Lencastre H, Oliveira D, Tomasz A. Antibiotic resistant *Staphylococcus aureus*: a paradigm of adaptive power. *Curr Opin Microbiol* 2007; 10:428–435.
- De Vlieghe S, Laevens H, Devriese LA, et al. Parturition teat apex colonization with *Staphylococcus chromogenes* in dairy heifers is associated with low somatic cell count in early lactation. *Vet Microbiol* 2003; 92:245–252.
- Devriese LA, Schleifer KH, Adegoke GO. Identification of coagulase-negative staphylococci from farm animals. *J Appl Bacteriol* 1983; 58:45–55.
- García-Álvarez L, Holden MT, Lindsay H, et al. Methicillin-resistant *Staphylococcus aureus* with a novel *mecA* homologue in human and bovine populations in the UK and Denmark: a descriptive study. *Lancet Infect Dis* 2011; 11:595–603.
- Gloor S, Bontadina F, Hegglin D, et al. The rise of urban fox populations in Switzerland. *Mamm Biol* 2001; 66:155–164.
- Hariharan H, Matthew V, Fountain J, et al. Aerobic bacteria from mucous membranes, ear canals, and skin wounds of feral cats in Grenada, and the antimicrobial drug susceptibility of major isolates. *Comp Immunol Microbiol Infect Dis* 2011; 34: 129–134.
- Hauschild T. Phenotypic and genotypic identification of staphylococci isolated from wild small mammals. *Syst Appl Microbiol* 2001; 24:411–416.

- Heuer H, Schmitt H, Smalla K. Antibiotic resistance gene spread due to manure application on agricultural fields. *Curr Opin Microbiol* 2011; 14:236–243.
- Hope R, Livermore DM, Brick G, et al. BSAC Working Parties on Resistance Surveillance. Non-susceptibility trends among staphylococci from bacteraemias in the UK and Ireland, 2001–06. *J Antimicrob Chemother* 2008; 62(Suppl 2):ii65–74.
- Irlinger F. Safety assessment of dairy microorganisms: Coagulase-negative staphylococci. *Int J Food Microbiol* 2008; 126: 302–310.
- Kawano J, Shimizu A, Saitoh Y, et al. Isolation of methicillin-resistant coagulase-negative staphylococci from chickens. *J Clin Microbiol* 1996; 34:2072–2077.
- Kirby WMM. Extraction of a highly potent penicillin inactivator from penicillin resistant staphylococci. *Science* 1944; 99:452–453.
- Kloos WE, Zimmerman RJ, Smith RF. Preliminary studies on the characterization and distribution of *Staphylococcus* and *Micrococcus* species on animal skin. *Appl Environ Microbiol* 1976; 31:53–59.
- Kozak GK, Boerlin P, Janecko N, et al. Antimicrobial resistance in *Escherichia coli* isolates from swine and wild small mammals in the proximity of swine farms and in natural environments in Ontario, Canada. *Appl Environ Microbiol* 2009; 75:559–566.
- Literak I, Dolejska M, Radimersky R, et al. Antimicrobial-resistant faecal *Escherichia coli* in wild mammals in central Europe: multiresistant *Escherichia coli* producing extended-spectrum beta-lactamases in wild boars. *J Appl Microbiol* 2009; 108:1702–1711.
- Loeffler A, Lloyd DH. Companion animals: a reservoir for methicillin-resistant *Staphylococcus aureus* in the community? *Epidemiol Infect* 2010; 138:595–605.
- Loeffler A, Pfeiffer DU, Lindsay JA, et al. Prevalence of and risk factors for MRSA carriage in companion animals: a survey of dogs, cats and horses. *Epidemiol Infect* 2010; 14:1–10.
- Martínez JL. Antibiotics and antibiotic resistance genes in natural environments. *Science* 2008; 321:365–367.
- Merlino J, Watson J, Rose B, et al. Detection and expression of methicillin/oxacillin resistance in multidrug-resistant and non-multidrug-resistant *Staphylococcus aureus* in Central Sydney, Australia. *J Antimicrob Chemother* 2002; 49:793–801.
- Millán J, Jiménez MA, Viota M, et al. Disseminated bovine tuberculosis in a wild red fox (*Vulpes vulpes*) in southern Spain. *J Wildl Dis* 2008; 44:701–706.
- Moodley A, Guardabassi L. Clonal spread of methicillin-resistant coagulase-negative staphylococci among horses, personnel and environmental sites at equine facilities. *Vet Microbiol* 2009; 137:397–401.
- Nagase N, Sasaki A, Yamashita K, et al. Isolation and species distribution of staphylococci from animal and human skin. *J Vet Med Sci* 2002; 64:245–250.
- Olsson-Liljequist B, Köljalg S, Karlsson I, et al. Calibration of fusidic acid disk diffusion susceptibility testing of *Staphylococcus aureus*. *Acta Path Micro Im B* 2002; 110:690–696.
- Owens CD, Stoessel K. Surgical site infections: epidemiology, microbiology and prevention. Review. *J Hosp Infect* 2008; 70(Suppl 2):3–10.
- Patel A, Lloyd DH, Lamport AI. Antimicrobial resistance of feline staphylococci in south-eastern England. *Vet Derm* 1999; 10:257–261.
- Piette A, Verschraegen G. Role of coagulase-negative staphylococci in human disease. *Vet Microbiol* 2009; 134:45–54.
- Rolland RM, Hausfater F, Marshall B, et al. Antibiotic-resistant bacteria in wild primates: increased prevalence in baboons feeding on human refuse. *Appl Environ Microbiol* 1985; 49:791–794.
- Schleifer KH, Kilpper-Baelz R, Devriese LA. *Staphylococcus arlettae* sp. nov., *S. equorum* sp. nov. and *S. kloosii* sp. nov.: Three new coagulase-negative, novobiocin-resistant species from animals. *Syst Appl Microbiol* 1984; 5:501–509.
- Scott E, Duty S, Callahan M. A pilot study to isolate *Staphylococcus aureus* and methicillin-resistant *S. aureus* from environmental surfaces in the home. *Am J Infect Control* 2008; 36:458–460.
- Soge OO, Meschke JS, No DB, et al. Characterization of methicillin-resistant *Staphylococcus aureus* and methicillin-resistant coagulase-negative *Staphylococcus* spp. isolated from US West Coast public marine beaches. *J Antimicrob Chemother* 2009; 64:1148–1155.
- Stepanovic S, Dakic I, Morrison D, et al. Identification and characterization of clinical isolates of members of the *Staphylococcus sciuri* group. *J Clin Microbiol* 2005; 43:956–958.
- Tacconelli E. Antimicrobial use: risk driver of multidrug resistant microorganisms in healthcare settings. *Curr Opin Infect Dis* 2009; 22:352–358.
- Takahashi T, Satoh I, Kikuchi N. Phylogenetic relationships of 38 taxa of the genus *Staphylococcus* based on 16S rRNA gene sequence analysis. *Int J Syst Bacteriol* 1999; 49:725–728.
- The Clinical and Laboratory Standards Institute (CLSI). Performance standards for antimicrobial disk and dilution susceptibility test for bacteria isolates from animals: Information Supplement. CLSI approved standard M31-S1 vol. 24. 2004. Clinical and Laboratory Standards Institute, Wayne, PA.
- Van de Giessen AW, van Santen-Verheuveld PD, Hengeveld TB, et al. Occurrence of methicillin-resistant *Staphylococcus aureus* in rats living on pig farms. *Prev Vet Med* 2009; 91: 270–273.
- Van Duijkeren E, Box ATA, Heck MEOC, et al. Methicillin-resistant staphylococci isolated from animals. *Vet Microbiol* 2004;103:91–97.
- Voss A, Loeffen F, Bakker J, et al. Methicillin-resistant *Staphylococcus aureus* in pig farming. *Emerg Infect Dis* 2005; 11:1965–1966.
- Wu SW, de Lencastre H, Tomasz A. Recruitment of the *mecA* gene homologue of *Staphylococcus sciuri* into a resistance determinant and expression of the resistant phenotype in *Staphylococcus aureus*. *J Bacteriol* 2001; 183:2417–2424.
- Zadoks RN, Watts JL. Species identification of coagulase-negative staphylococci: Genotyping is superior to phenotyping. *Vet Microbiol* 2009; 134:20–28.
- Zhang Y, Agidi S, Le Jeune JT. Diversity of staphylococcal cassette chromosome in coagulase-negative staphylococci from animal sources. *J Appl Microbiol* 2009; 107:1375–1383.

Address correspondence to:

Anette Loeffler  
 Department of Veterinary Clinical Sciences  
 Royal Veterinary College  
 Hawkshead Lane, Hatfield  
 North Mymms, Hertfordshire AL9 7TA  
 United Kingdom

E-mail: aloeffler@rvc.ac.uk

**This article has been cited by:**

1. Katarzyna Garbacz, Sabina #arnowska, Lidia Piechowicz, Krystyna Haras. 2012. Staphylococci Isolated from Carriage Sites and Infected Sites of Dogs as a Reservoir of Multidrug Resistance and Methicillin Resistance. *Current Microbiology* .  
[CrossRef]