



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Covid-19 Obesity, Deprivation and Death

Citation for published version:

Simpson, AHRW, Simpson, CJ, Frost, H & Welburn, SC 2020, 'Covid-19 Obesity, Deprivation and Death', *Journal of Global Health*. <https://doi.org/10:020389>

Digital Object Identifier (DOI):

[10:020389](https://doi.org/10:020389)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Global Health

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Cite as: Simpson HRW, Simpson CJ, Frost F, Welburn SC. COVID-19: Obesity, deprivation and death. J Glob Health 2020;10:020389.

COVID-19: Obesity, deprivation and death

Hamish RW Simpson, Cameron J Simpson, Helen Frost, Susan C Welburn

¹Edinburgh Medical School, University of Edinburgh, Edinburgh Scotland

²NHS Scotland, Edinburgh, Scotland

³Edinburgh Napier University, Edinburgh, Scotland

⁴Infection Medicine, College of Medicine and Veterinary Medicine, University of Edinburgh, Edinburgh, Scotland

⁵Zhejiang University-University of Edinburgh Institute, Zhejiang University School of Medicine, Zhejiang, PR China

Photo: From Hamish Simpson's own collection, used with permission.



The Covid-19 has disrupted health systems, even within the most developed economies, with case fatality rates in certain countries exceeding 10% [1].

COVID-19 deaths have been linked to diabetes and hypertension [2], which also correlate with obesity. As adipose tissue is prone to infection by the virus and produces high levels of inflammatory

cytokines [3], it may feed the cytokine storm [4] suffered by many COVID-19 fatalities suggesting obesity could be the key factor.

COVID-19 nation death rates vary greatly, but countries with low obesity rates have consistently low fatality rates; specifically, Thailand, China, South Korea, Japan, Pakistan, Philippines and India are all countries of over 10 million population, but with a national obesity rate of 10% or less [5], which all have less than 50 deaths per million [1]. There are clearly a multitude of factors that contribute to the national death rates. However, countries with obesity rates of over 20% have far more variable mortality rates.

The poorest populations are most at risk of chronic conditions, which may put them at risk of dying from COVID19 [5], yet war torn countries such as Syria have not had the exponential increases in deaths observed in Europe and North America; suggesting that when comparing the effect on mortality, of different national strategies for COVID-19, national obesity rates should be considered. National mitigation measures including lockdowns and social distancing may have to extend beyond 2020/21. During this time, individuals have been advised to do mild-to-moderate exercise as it can be of benefit in fighting viral diseases [6]. In addition, individuals should be advised to lose weight (with a mild caloric restriction) to minimise the risk of succumbing to COVID-19.

The pressing need for this is illustrated by the analysis of risk factors for critical illness; patients with a BMI >40 had a far higher risk (odds ratio OR 6.2) of requiring hospitalisation [7]. As over one-third of Americans, over one-fifth of Europeans and 13% of the adults in the world are obese [5], a very high percent of the population is at risk of severe symptomatology.

Obesity rates rise with economic development of a country [8]. Paradoxically, the relationship of socioeconomic status (SES) to obesity changes, those of higher SES tend to be obese in lower income countries, whereas those of lower SES tend to be obese in high income countries. Thus, poorer individuals in wealthier nations are at particularly high risk of dying from COVID-19, yet due to inequalities, they will have most difficulty in accessing essential life care support. Unfortunately, interventions to prevent transmission have a particularly negative impact on these already

impoverished communities. Ethnic minorities have been particularly hard hit by COVID19, highlighting racial as well as economic inequalities [9].

With COVID-19, economies have been pulled into recession and unemployment rates have spiralled, exhausting national welfare safety nets and placing the burden onto the individual. Yet, those most at risk from this disease are the poorest in society, who have been most impacted by the mitigation strategies and those least able to access healthcare. Global health promotion strategies should emphasise the benefits of physical activity and weight loss interventions (such as those described by Brown et al [10]) in the fight against this pandemic, and country specific policies need to be developed quickly to support people who are most at risk.

Acknowledgements: None.

Funding: None received.

Authorship contributions: all authors contributed to the literature search, data review and writing of the manuscript

Competing interests: The authors completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available upon request from the corresponding author) and declare no conflicts of interest.

REFERENCES

- 1 Reported Cases and Deaths by Country, Territory, or Conveyance. Available: https://www.worldometers.info/coronavirus/?utm_campaign=homeAdvegas1?#countries. Accessed: Sept 2020
- 2 Guan WJ et al. Comorbidity and its impact on 1590 patients with Covid-19 in China: A Nationwide Analysis. *Eur Respir J.* 2020;55:2000547.
- 3 Fain JN. Release of interleukins and other inflammatory cytokines by human adipose tissue is enhanced in obesity and primarily due to the non-fat cells. *Vitam Horm.* 2006;74:2006.
- 4 Mehta P, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ, on behalf of the HLH. Across Speciality Collaboration, UK. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet.* 2020;395:10229:1033-34.

- 5 Ahmed F, Ahmed N, Pissarides C, Stiglitz J. Why inequality could spread COVID-19. *Lancet Public Health*. 2020;5:e240.
- 6 Luzi L, Radaelli MG. Influenza and obesity: its odd relationship and the lessons for COVID-19 pandemic. *Acta Diabetol*. 2020;57:759-64.
- 7 Petrilli CM, Jones SA, Yang J, Rajagopalan H, O'Donnell LF, Chernyak Y, et al. Factors associated with hospitalization and critical illness among 4,103 patients with COVID-19 disease in New York City. *medRxIV*. 2020. DOI: <https://doi.org/10.1101/2020.04.08.20057794>.
- 8 Pampel FC, Denney JT, Krueger PM. Obesity, SES, and economic development: A test of the reversal hypothesis. *Soc Sci Med*. 2012;74:1073-81.
- 9 Garg S, Kim L, Whitaker M, O'Halloran A, Cummings C, Holstein R et al. Hospitalization rates and characteristics of patients hospitalized with laboratory-confirmed coronavirus disease 2019 — COVID-NET, 14 States, March 1–30, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69:458–464.
- 10 Brown A, Dornhorst A, McGowan B, Omar O, Leeds AR, Taheri S, et al. Low-energy total diet replacement intervention in patients with type 2 diabetes mellitus and obesity treated with insulin: a randomized trial. *BMJ Open Diabetes Res Care*. 2020;8:e001012.

Correspondence to:

Prof Hamish Simpson
49 Little France Crescent
SU.303, Chancellor's Building
Edinburgh EH16 4SB
Scotland, UK
Hamish.simpson@ed.ac.uk