



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## An alternative route to the system-size expansion

**Citation for published version:**

Cianci, C, Schnoerr, D, Piehler, A & Grima, R 2017, 'An alternative route to the system-size expansion' Journal of Physics A: Mathematical and Theoretical, vol. 50, no. 39, 395003. DOI: 10.1088/1751-8121/aa85aa

**Digital Object Identifier (DOI):**

[10.1088/1751-8121/aa85aa](https://doi.org/10.1088/1751-8121/aa85aa)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Journal of Physics A: Mathematical and Theoretical

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# An alternative route to the system-size expansion

Claudia Cianci, David Schnoerr, Andreas Piehler, Ramon Grima

*School of Biological Sciences, University of Edinburgh, UK*

## Abstract

The master equation is rarely exactly solvable and hence various means of approximation have been devised. A popular systematic approximation method is the system-size expansion which approximates the master equation by a generalised Fokker-Planck equation. Here we first review the use of the expansion by applying it to a simple chemical system. The example shows that the solution of the generalised Fokker-Planck equation obtained from the expansion is generally not positive definite and hence cannot be interpreted as a probability density function. Based on this observation, one may also *a priori* conclude that moments calculated from the solution of the generalised Fokker-Planck equation are not accurate; however calculation shows these moments to be in good agreement with those obtained from the exact solution of the master equation. We present an alternative simpler derivation which directly leads to the same moments as the system-size expansion but which bypasses the use of generalised Fokker-Planck equations, thus circumventing the problem with the probabilistic interpretation of the solution of these equations.

## I. INTRODUCTION

The Markovian description of stochastic systems with discrete state space is generally described by means of a master equation [1]. The solution of the latter gives the probability that the system of interest is in a given configuration at a given time. However, exact solution of the master equation is rarely possible because of the typically large (or infinite) dimensionality of the state space. The most common strategy to bypass this difficulty involves the use of stochastic simulations via the stochastic simulation algorithm (SSA) [2] or one of its several variants. However, such simulations can easily become computationally expensive when modelling realistic systems, thus making analytical approximation procedures an appealing alternative.

One of the most well-known approximation methods is van Kampen's system-size expansion (SSE) [1], a perturbative expansion of the master equation in the inverse system size, for systems which are deterministically monostable. In the limit of infinitely large system sizes, consideration of the leading order terms of the SSE shows that the mean concentrations of the master equation agree with those of the deterministic rate equations, while the distribution of fluctuations about the mean is Gaussian and given by a Fokker-Planck equation with linear drift and diffusion coefficients. The latter is often referred to as the "linear-noise approximation" (LNA) and is widely used in the literature (see for example [1, 3–5]). Consideration of higher orders of the expansion lead to generalised Fokker-Planck equations (GFPE) with third or higher order derivatives [6]. These higher-order terms have recently been used [7–11] to compute corrections to the mean concentration solution of the rate equations and to the second moments given by the LNA (for a review of these results see [12]). These corrections are applicable when the system size is of intermediate size. Although they are computationally advantageous and typically highly accurate when compared to stochastic simulations [13–15], the GFPE from which they are computed has been subject to strong criticisms. Pawula's theorem [16, 17] states that the solution of the GFPE is a positive quantity only if the GFPE has at most second-order derivatives or else if the GFPE has an infinite number of terms. In other words, while the solution of the Fokker-Planck equation associated with the LNA can be interpreted as a probability density function, the solution of the GFPE with third or higher derivatives can not be so interpreted. Thus the moments computed from such an equation are dubious. Based on such arguments one may dismiss

any SSE computations beyond the LNA [18]. Hence the dilemma: why are the moments calculated using higher-order terms beyond the LNA quantitatively accurate despite being derived from a GFPE which has no apparent meaningful probabilistic interpretation?

In this paper, we propose a novel method, which leads to the same moment equations as obtained using the GFPE with third and higher-order derivatives derived from the SSE, but which avoids the problems outlined above. In particular, the method relies on an expansion of the moment equations directly obtained from the master equation and hence bypasses the use of a time-evolution equation for the approximate probability density function. To be specific, we shall consider a general chemical reaction system containing different species interacting via a number of chemical reactions in a well-mixed volume. In this context the master equation is known as the “chemical master equation” (CME) and the system size corresponds to the volume of the compartment in which the reactions occur. We emphasise however that all results derived in this paper also apply to master equations that allow a SSE but are not of CME type (see for example [19, 20]).

This paper is structured as follows. In Section II we introduce a simple, instructive one variable example and show that while the solution of the corresponding GFPE with up to fourth- and sixth-order derivatives can become negative, the moments calculated from such a solution are highly accurate, thereby illustrating the aforementioned dilemma. In Section III, we present the general derivation of the SSE leading to a GFPE with up to fourth-order derivatives for a general multi-species chemical system. In Section IV we present our new expansion method starting directly from the moment equations of the CME and which bypass the use of any type of GFPE. Subsequently in Section V we show that the equations from the conventional SSE derivation of Section III match exactly the equations derived from the alternative method in Section IV. Finally, we briefly summarize our results and conclude in Section VI.

## II. AN ILLUSTRATIVE ONE SPECIES EXAMPLE

In order to demonstrate the probabilistic interpretation problem of GFPEs, we consider the following simple chemical reaction system:



The species  $X$  gets created at rate  $k_1$  and irreversibly forms dimers with rate  $k_2$  (which we do not follow and hence we do not label). The corresponding CME is given by:

$$\partial_t P(n, t) = \Omega \left[ (E^{-1} - 1) k_1 + (E^2 - 1) \frac{k_2}{\Omega^2} n(n-1) \right] P(n, t), \quad (2)$$

where  $P(n, t)$  is the probability of  $n$  particles of  $X$  being in the system at time  $t$ ,  $E^{\pm x}$  are the step operators that replace  $n$  with  $n \pm x$ , i.e.,  $E^{\pm x} f(n) = f(n \pm x)$  for a function  $f(n)$ , and  $\Omega$  is the volume of the compartment. The SSE makes the ansatz [1]:

$$\frac{n}{\Omega} = \phi + \frac{\epsilon}{\sqrt{\Omega}}, \quad (3)$$

where  $\phi$  is the solution of the deterministic rate equations. We thus divide the concentration  $n/\Omega$  into a deterministic contribution  $\phi$  and a contribution  $\epsilon$  which provides corrections to the latter. The corresponding deterministic rate equations are given by:

$$\partial_t \phi = k_1 - 2k_2 \phi^2. \quad (4)$$

The ansatz in (3) effectively leads to a change of variables from  $n$  to  $\epsilon$ . We thus have to transform from the distribution  $P(n, t)$  over  $n$  to a new distribution  $\Pi(\epsilon, t)$  over  $\epsilon$ . Performing the transformation in variables on the CME (2) and taking the limit of large  $\Omega$  (see [8] for technical details of the general procedure), one finds that the CME in Eq. (2) can be approximated by the GFPE:

$$\begin{aligned} \frac{\partial \Pi(\epsilon, t)}{\partial t} = & \Omega^0 \left[ \frac{1}{2} (k_1 + 4k_2 \phi^2) \frac{\partial^2 \Pi(\epsilon, t)}{\partial \epsilon^2} + 4k_2 \phi \frac{\partial}{\partial \epsilon} (\epsilon \Pi(\epsilon, t)) \right] \\ & + \Omega^{-1/2} \left[ -\frac{1}{6} (k_1 - 8k_2 \phi^2) \frac{\partial^3 \Pi(\epsilon, t)}{\partial \epsilon^3} + 4k_2 \phi \frac{\partial^2}{\partial \epsilon^2} (\epsilon \Pi(\epsilon, t)) \right. \\ & \left. - 2k_2 \phi \frac{\partial \Pi(\epsilon, t)}{\partial \epsilon} + 2k_2 \frac{\partial}{\partial \epsilon} (\epsilon^2 \Pi(\epsilon, t)) \right] \\ & + \Omega^{-1} \left[ \frac{1}{24} (k_1 + 2^4 k_2 \phi^2) \frac{\partial^4 \Pi(\epsilon, t)}{\partial \epsilon^4} + \frac{8}{3} k_2 \phi \frac{\partial^3 (\epsilon \Pi(\epsilon, t))}{\partial \epsilon^3} - 2k_2 \frac{\partial^2 \Pi(\epsilon, t)}{\partial \epsilon^2} \phi \right. \\ & \left. + 2k_2 \frac{\partial^2}{\partial \epsilon^2} (\epsilon^2 \Pi(\epsilon, t)) - 2k_2 \frac{\partial}{\partial \epsilon} (\epsilon \Pi(\epsilon, t)) \right] \\ & + \Omega^{-3/2} \left[ -2k_2 \frac{\partial^2}{\partial \epsilon^2} (\epsilon \Pi(\epsilon, t)) + \frac{4}{3} k_2 \frac{\partial^3}{\partial \epsilon^3} (\epsilon^2 \Pi(\epsilon, t)) - \frac{4}{3} k_2 \phi \frac{\partial^3}{\partial \epsilon^3} (\Pi(\epsilon, t)) \right. \\ & \left. + \frac{4k_2 \phi}{3} \frac{\partial^4}{\partial \epsilon^4} (\epsilon \Pi(\epsilon, t)) - \frac{1}{120} (k_1 + (-2)^5 k_2 \phi^2) \frac{\partial^5}{\partial \epsilon^5} \Pi(\epsilon, t) \right] \\ & + \Omega^{-2} \left[ \frac{1}{6!} (k_1 + 2^6 k_2 \phi^2) \frac{\partial^6}{\partial \epsilon^6} \Pi(\epsilon, t) + \frac{1}{5!} 2^6 k_2 \phi \frac{\partial^5}{\partial \epsilon^5} (\epsilon \Pi(\epsilon, t)) \right. \\ & \left. + \frac{1}{4!} 2^4 k_2 \frac{\partial^4}{\partial \epsilon^4} ((\epsilon^2 - \phi) \Pi(\epsilon, t)) - \frac{1}{3!} 2^3 k_2 \frac{\partial^3}{\partial \epsilon^3} (\epsilon \Pi(\epsilon, t)) \right] + O(\Omega^{-5/2}). \quad (5) \end{aligned}$$




Figure 1: The steady-state solution of the GFPE in (6) obtained by truncating van Kampen's SSE to orders  $\Omega^0$  (red, LNA),  $\Omega^{-1}$  (blue) and  $\Omega^{-2}$  (orange) for the dimerization reaction system in (1). The grey dots show the exact steady-state solution of the CME. The parameters used are  $k_1 = k_2 = 1$  and  $\Omega = 1$ . Note that while the LNA solution is positive for all  $n$ , the higher order solutions (blue, orange) are negative for some values of  $n$ . Hence while the LNA has a probabilistic interpretation, the higher-order GFPE solutions do not.

Note that truncating terms on the right hand side of the above equation to order  $\Omega^0$  leads to the LNA, a Fokker-Planck equation with a drift term linear in  $\epsilon$  and a diffusion coefficient which is independent of  $\epsilon$ . This equation admits a Gaussian solution for the distribution of fluctuations about the solution of the rate equations. A closed formula for the non-Gaussian solution of the GFPE for any deterministically monostable single species system, truncated to any order, has been derived recently [14]. Applying this general formula, we find that the steady-state solution of the GFPE in Eq. (5) is given by:

$$\Pi(\epsilon) = \pi_0(\epsilon) \left( 1 + \sum_{j=1}^4 \Omega^{-j/2} \sum_{m=1}^{3j} a_m^{(j)} \psi_m(\epsilon) \right) + O(\Omega^{-5/2}). \quad (6)$$

The first term  $\pi_0(\epsilon)$  is the Gaussian solution of the LNA and reads:

$$\pi_0(\epsilon) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{\epsilon^2}{2\sigma^2}\right), \quad (7)$$

where  $\sigma^2 = \frac{3}{4}\phi_s$  and  $\phi_s = \sqrt{k_1/2k_2}$ . The latter is the steady-state solution of the rate equation in (4). The corrections beyond the LNA in (6) are given in terms of the functions  $\psi_m(\epsilon)$

which are proportional to Hermite polynomials  $H_m$  as follows:

$$\psi_m(\epsilon) = \frac{1}{\sigma^m} H_m\left(\frac{\epsilon}{\sigma}\right). \quad (8)$$

The non-vanishing coefficients  $a_m^{(i)}$  in Eq. (6) read:

$$\begin{aligned} a_1^{(1)} &= \frac{1}{8}, a_3^{(1)} = \frac{7}{96}\phi_s, \\ a_2^{(2)} &= \frac{5}{128}, a_4^{(2)} = \frac{23}{1536}\phi_s, a_6^{(2)} = \frac{49}{18432}\phi_s^2, \\ a_1^{(3)} &= \frac{3}{128\phi_s}, a_3^{(3)} = \frac{13}{3072}, a_5^{(3)} = \frac{37}{10240}\phi_s, a_7^{(3)} = \frac{7}{9216}\phi_s, a_9^{(3)} = \frac{343}{5308416}\phi_s^{3/2}, \\ a_2^{(4)} &= \frac{3}{2048\phi_s}, a_4^{(4)} = -\frac{13}{98304}, a_6^{(4)} = \frac{2089}{2949120}\phi_s, a_8^{(4)} = \frac{4171}{23592960}\phi_s^2, a_{10}^{(4)} = \frac{2009}{84934656}\phi_s^{3/2}, a_{12}^{(4)} = \frac{2401}{2038431744}\phi_s^4. \end{aligned} \quad (9)$$

Fig. 1 shows the solutions to the GFPE given in Eq. (6) truncated to orders  $\Omega^0$  (red),  $\Omega^{-1}$  (blue) and  $\Omega^{-2}$  (orange). The latter correspond to the GFPE with at most second-, fourth- and sixth-order derivatives, respectively. Note that Fig. 1 shows distributions in the variable  $n$ , which is obtained by reversing the transformation of variables from distributions in  $\epsilon$  to distributions over the variable  $n$ .

Note that truncating the GFPE to higher-order leads to a better agreement of the solution of the GFPE with that of the exact solution of the CME for positive  $n$ . However this is achieved at the expense of negative probabilities for some negative  $n$  values (for other examples, such as the simple birth-death process, the higher-order GFPE solution leads to negative probabilities also for positive  $n$  values – see Fig. 1a in [14]). Note that all GFPE's (including the LNA) are not just defined for positive molecule numbers  $n$  but also for negative ones. This is in contrast to the CME which only allows transitions between positive molecule numbers. However for the LNA this is not a major cause of concern since the probability is positive and hence the LNA constitutes a stochastic process over the real domain; in contrast, due to negative probabilities, the higher-order GFPE's do not allow a probabilistic interpretation. Curiously, however, the means and variances of the continuous GFPE distributions computed over the open interval  $n = (-\infty, \infty)$  are found to progressively approach the exact CME value as we include higher-order derivatives (see Table I). Hence we have a dilemma: *despite being derived from a distribution that is not positive definite, these moments are closer to the true moments of the CME, when compared to the moments calculated from the positive-definite distribution of the LNA.*

	Exact	$\Omega^0$	$\Omega^{-1}$	$\Omega^{-2}$
mean	0.888	0.707	0.832	0.865
variance	0.600	0.530	0.593	0.601


Table I: Comparison of the mean and variance in molecule numbers calculated from the exact CME distribution and the solutions of the GFPE truncated to orders  $\Omega^0$  (LNA),  $\Omega^{-1}$  and  $\Omega^{-2}$  (the orders of  $\Omega^{-1/2}$  and  $\Omega^{-3/2}$  do not contribute to the moments). Parameters are as in Fig. 1 where we also show the corresponding distributions. Note that the mean and variance become more accurate with increasing order. The exact values are computed using formulae (68)-(69) in Reference [7].

More generally one finds that for a fixed value of the system size  $\Omega$ , consideration of higher-order derivatives of the GFPE leads to more accurate estimates of the moments until a certain order, after which the accuracy rapidly decreases; this threshold order increases with the system size and hence strictly speaking, the expansion is valid to all orders only in the limit  $\Omega \rightarrow \infty$  [21]. This also implies that the SSE is divergent for finite  $\Omega$ . This is typical of series derived from perturbation methods [22]. It is not possible to *a priori* know the threshold order however generally if the addition of a new term leads to either a large relative change in the value of a moment or if it leads to a negative value of the average of the molecule numbers raised to any power then clearly one should not add further terms to the series.

The accuracy of moments computed from higher order truncations of the GFPE (below the threshold order) leads one to surmise that it may be possible to derive them using an approach which bypasses the use of the GFPE and hence the knotty issue of negative probabilities. The rest of this paper is dedicated to derive such an approach.

### III. THE SYSTEM-SIZE EXPANSION FOR A GENERAL CHEMICAL SYSTEM

We start by reviewing the derivation of the SSE to higher orders introduced in [7]. Consider a system of  $N$  chemical species that interact through a set of  $R$  chemical reactions where the  $r^{th}$  reaction has the form:





Here the index  $r$  takes values between 1 and  $R$ , and  $X_i$  denotes species  $i$ ,  $i = 1, \dots, N$ . The integers  $s_{ir}$  and  $h_{ir}$  are the stoichiometric coefficients and  $k_r$  is the macroscopic reaction rate. We can write the corresponding CME in compact form using step operators as [1]:

$$\partial_t P(\vec{n}, t) = \Omega \sum_{r=1}^R \left( \prod_{i=1}^N E_i^{-S_{ir}} - 1 \right) f_r(\vec{n}, \Omega) P(\vec{n}, t), \quad (11)$$

where  $\vec{n} = (n_1, \dots, n_N)$ ,  $n_i$  is the molecule number of species  $X_i$ ,  $E_i^x$  is a step operator returning  $f_r(\vec{n}, \Omega) P(\vec{n}, t)$  but with  $n_i$  replaced by  $n_i + x$ , and we have defined the stoichiometric matrix  $S_{ir} = h_{ir} - s_{ir}$ . The microscopic rate function  $f_r(\vec{n}, \Omega)$  depends on the type of the  $r^{\text{th}}$  chemical reaction. The probability that a reaction occurs in a time interval  $[t, t + dt)$  somewhere in the volume  $\Omega$  is given by  $\Omega f_r(\vec{n}, \Omega) dt$ .

We assume mass action kinetics, and specifically we consider reactions involving at most two reactant molecules, i.e.,  $\sum_{i=1}^N s_{ir} \leq 2 \forall r$ , since reactions involving three or more molecules are relatively rare. This means that we can write the functions  $f_r(\vec{n}, t)$  in (11) in the following form:

$$\Omega f_r(\vec{n}, \Omega) = \left[ \Delta_r \Omega + \sum_{k'=1}^N \beta_r^{k'} n_{k'} + \sum_{k', s=1, s \neq k'}^N \frac{\gamma_r^{k's}}{\Omega} n_{k'} n_s + \sum_{k'=1}^N \frac{\zeta_r^{k'k'}}{\Omega} n_{k'} (n_{k'} - 1) \right], \quad (12)$$

where we explicitly split up the term into contributions from different types of possible reactions. The first and second term represent zeroth and first order (unimolecular) reactions, respectively. The third term corresponds to a second order (bimolecular) reaction, where the two reactant molecules are of different type, while the last term corresponds to a bimolecular reaction involving two identical reactant molecules. Note that for a given reaction  $r$ , only one of the coefficients  $\Delta_r, \beta_r^{k'}, \gamma_r^{k's}$  and  $\zeta_r^{k'k'}$  in Eq. (12) is non-zero, depending on the type of reaction, and it is equal to  $k_r$ .

To derive the SSE equations one makes the following ansatz:

$$\frac{n_i}{\Omega} = \phi_i + \frac{\epsilon_i}{\sqrt{\Omega}}, \quad i = 1, \dots, N. \quad (13)$$

Note that this is the same ansatz that we made in Eq. (3) for the single species system, but this time it is applied to the concentrations of each species. The  $\phi_i$  are the solution of the deterministic rate equations. Next, we transform the CME in Eq. (11) from the variables  $\vec{n}$  to the variables  $\vec{\epsilon}$ , leading to a GFPE for the distribution  $\Pi(\vec{\epsilon}, t)$ . In order to perform this transformation, we transform the time derivative, expand the step operators  $E_i^x$  in powers

of  $\Omega^{-1/2}$  (for the explicit expression see [7]) and transform the functions  $f_r$  accordingly. This leads to:

$$\frac{\partial \Pi(\vec{\epsilon}, t)}{\partial t} - \Omega^{1/2} \sum_{i=1}^N \frac{d\phi_i}{dt} \frac{\partial \Pi(\vec{\epsilon}, t)}{\partial \epsilon_i} = -\Omega^{1/2} \sum_{i=1}^N \sum_{r=1}^R S_{ir} f_r(\vec{\phi}) \frac{\partial \Pi(\vec{\epsilon}, t)}{\partial \epsilon_i} + \sum_{i=0}^{\infty} \Omega^{-i/2} L^{(i)} \Pi(\vec{\epsilon}, t), \quad (14)$$

where  $L^{(i)}$  are differential operators – the reader is referred to [6] for their explicit definitions.

Now by the law of mass action, one can deduce that the deterministic rate equations for the system (10) are:

$$\begin{aligned} \frac{d\phi_i}{dt} &= \sum_{r=1}^R S_{ir} f_r(\vec{\phi}), \\ f_r(\vec{\phi}) &= \lim_{\Omega \rightarrow \infty} f_r(\vec{n} = \Omega \vec{\phi}, \Omega) = \Delta_r + \sum_{k'=1}^N \beta_r^{k'} \phi_{k'} + \sum_{k', s=1, s \neq k'}^N \gamma_r^{k' s} \phi_{k'} \phi_s + \sum_{k'=1}^N \zeta_r^{k' k'} \phi_{k'}^2, \end{aligned} \quad (15)$$

where  $\vec{\phi}$  is the vector of macroscopic concentrations and  $f_r(\vec{\phi})$  is the macroscopic rate function of the  $r^{th}$  reaction. In the macroscopic limit, terms of order  $\Omega^{1/2}$  in Eq. (14) dominate. However by the assumption that the CME must agree with the rate equations in the macroscopic limit, one finds that terms of order  $\Omega^{1/2}$  on both sides of the transformed CME disappear. Hence the transformed CME Eq. (14) reads:

$$\frac{\partial \Pi(\vec{\epsilon}, t)}{\partial t} = \left( \Omega^0 L^{(0)} + \Omega^{-1/2} L^{(1)} + \Omega^{-1} L^{(2)} \right) \Pi(\vec{\epsilon}, t) + O(\Omega^{-3/2}). \quad (16)$$

The form of the transformed CME in Eq. (16) suggests the perturbative solution:

$$\Pi(\vec{\epsilon}, t) = \sum_{j=0}^{\infty} \Pi_j(\vec{\epsilon}, t) \Omega^{-j/2}. \quad (17)$$

Substitution of this ansatz in Eq. (16) and collecting terms of equal orders leads to time-evolution equations for  $\Pi_j(\vec{\epsilon}, t)$ . From these equations we can subsequently obtain time-evolution equations for the pseudo-moments  $[\epsilon_k \epsilon_m \dots \epsilon_r]_j = \int d\vec{\epsilon} \epsilon_k \epsilon_m \dots \epsilon_r \Pi_j(\vec{\epsilon}, t)$ .

Finally the equations of the moments of the concentrations are obtained from the van

Kampen ansatz in Eq. (13) and are given by:

$$\left\langle \frac{n_i}{\Omega} \right\rangle = \phi_i + \frac{\langle \epsilon_i \rangle}{\sqrt{\Omega}} = \phi_i + \sum_{j=0}^3 [\epsilon_i]_j \Omega^{-(j+1)/2}, \quad (18)$$

$$\begin{aligned} \left\langle \frac{n_i n_l}{\Omega \Omega} \right\rangle &= \phi_i \phi_l + \frac{\phi_i \langle \epsilon_l \rangle + \phi_l \langle \epsilon_i \rangle}{\sqrt{\Omega}} + \frac{\langle \epsilon_i \epsilon_l \rangle}{\Omega} \\ &= \phi_i \phi_l + \Omega^{-1/2} (\phi_i [\epsilon_l]_0 + [\epsilon_i]_0 \phi_l) + \Omega^{-1} ([\epsilon_i \epsilon_l]_0 + \phi_i [\epsilon_l]_1 + \phi_l [\epsilon_i]_1) \\ &\quad + \Omega^{-3/2} ([\epsilon_i \epsilon_l]_1 + \phi_i [\epsilon_l]_2 + \phi_l [\epsilon_i]_2) + \Omega^{-2} ([\epsilon_i \epsilon_l]_2 + \phi_i [\epsilon_l]_3 + \phi_l [\epsilon_i]_3), \end{aligned} \quad (19)$$

$$\begin{aligned} \left\langle \frac{n_i n_k n_l}{\Omega^3} \right\rangle &= \phi_i \phi_k \phi_l + \Omega^{-1/2} \left( \phi_k \phi_l [\epsilon_i]_0 + (i \leftrightarrow k) + (k \leftrightarrow l) \right) + \Omega^{-1} \left( \phi_k [\epsilon_i \epsilon_l]_0 + (k \leftrightarrow i) + (i \leftrightarrow l) \right. \\ &\quad \left. + \phi_k \phi_l [\epsilon_i]_1 + (i \leftrightarrow k) + (k \leftrightarrow l) \right) + \Omega^{-3/2} \left( [\epsilon_k \epsilon_i \epsilon_l]_0 + \left[ \phi_k \phi_l [\epsilon_i]_2 + (i \leftrightarrow k) + (k \leftrightarrow l) \right. \right. \\ &\quad \left. \left. + \phi_k [\epsilon_i \epsilon_l]_1 + (i \leftrightarrow k) + (i \leftrightarrow l) \right] \right) + \Omega^{-2} \left( [\epsilon_k \epsilon_i \epsilon_l]_1 + \left[ \phi_k \phi_l [\epsilon_i]_3 + (i \leftrightarrow k) + (k \leftrightarrow l) \right. \right. \\ &\quad \left. \left. + \phi_k [\epsilon_i \epsilon_l]_2 + (i \leftrightarrow k) + (i \leftrightarrow l) \right] \right), \end{aligned} \quad (20)$$

where we truncated to order  $\Omega^{-2}$ . Note that the short hand notation  $(i \leftrightarrow k)$  stands for all the expressions of the same form as the one preceding the notation but with  $i$  and  $k$  interchanged. For example  $\phi_k \phi_l [\epsilon_i]_0 + (i \leftrightarrow k) + (k \leftrightarrow l)$  implies  $\phi_k \phi_l [\epsilon_i]_0 + \phi_i \phi_l [\epsilon_k]_0 + \phi_i \phi_k [\epsilon_l]_0$ .

Note that here we used the fact that:

$$\langle \epsilon_k \epsilon_m \dots \epsilon_r \rangle = \int d\vec{\epsilon} \epsilon_k \epsilon_m \dots \epsilon_r \Pi(\vec{\epsilon}, t) = \sum_{j=0}^{\infty} \Omega^{-j/2} [\epsilon_k \epsilon_m \dots \epsilon_r]_j, \quad (21)$$

which follows from Eq. (17) and the definition of the pseudo-moments given below it. This procedure can also be generalized to chemical systems with non-mass action kinetics and those reactions involving three or more reactant molecules [13].

In what follows, we shall use the Einstein summation convention where all twice repeated indices in product terms are understood to be summed over 1 to  $N$ . This notation will be used throughout the paper when it is necessary to write equations in compact form. To simplify the presentation, we define the following coefficients:

$$\begin{aligned} J_{ij\dots w}^{pt\dots z} &= \frac{\partial}{\partial \phi_p} \frac{\partial}{\partial \phi_t} \dots \frac{\partial}{\partial \phi_z} J_{ij\dots w}, \\ J_{ij\dots w} &= \sum_{r=1}^R S_{ir} S_{jr} \dots S_{wr} f_r(\vec{\phi}). \end{aligned} \quad (22)$$

Because of the quadratic form of  $f_r(\vec{\phi})$  (which stems from allowing at most bimolecular reactions), the only non-zero  $J$  coefficients are those involving at most second-order derivatives,

i.e., those with at most two superscripts. These are:

$$\begin{aligned}
J_{ij\dots w}^p &= \sum_{r=1}^R S_{ir} S_{jr} \dots S_{wr} \left[ \sum_{k'=1}^N \beta_r^{k'} \delta_{k',p} + \sum_{k',s=1,s \neq k'}^N \gamma_r^{k's} (\phi_s \delta_{k',p} + \phi_{k'} \delta_{s,p}) + \sum_{k'=1}^N 2\zeta_r^{k'k'} \phi_{k'} \delta_{k',p} \right], \\
J_{ij\dots w}^{pt} &= \sum_{r=1}^R S_{ir} S_{jr} \dots S_{wr} \left[ \sum_{k',s=1,s \neq k'}^N \gamma_r^{k's} (\delta_{k',p} \delta_{s,t} + \delta_{s,p} \delta_{k',t}) + \sum_{k'=1}^N 2\zeta_r^{k'k'} \delta_{k',p} \delta_{k',t} \right], \tag{23}
\end{aligned}$$

where  $\delta_{k',p}$  is the Kronecker delta.

The time-evolution equations for the pseudo-moments  $[\cdot]_j$  are needed to compute the moments in concentrations. These equations are obtained as detailed just before Eq. (18) and can be found in Appendix A.

In summary, the estimates of the first, second and third moments of the concentrations up to order  $\Omega^{-2}$  according to the SSE are generally given by Eqs. (18)-(20) together with the solution of the closed set of differential equations given by Eqs. (A1)-(A10). Equations for the fourth and higher-order moments in the concentrations can be derived similarly. For more details on the derivation we refer the reader to [7, 13].

#### IV. AN ALTERNATIVE ROUTE

As we saw in the last section, the final output of the SSE as given by Eqs. (18)-(20) is a series expansion of the moments in powers of  $\Omega^{-1/2}$ . The open question is whether this expansion is accurate given that it is derived from the GFPE in Eq. (16) which has higher than second-order derivatives and hence admits no probabilistic interpretation according to Pawula's theorem. To solve this apparent issue, we next formulate an alternative derivation which relies on a direct expansion of the moment equations obtained from the CME in the inverse system size, thereby avoiding the use of a GFPE. The procedure is as follows.

The moment equations are obtained by multiplying the CME in Eq. (11) on both sides with  $n_i \dots n_l$  and subsequently summing over all allowed values of the molecule numbers.

The moment equations of the first four moments read:

$$\partial_t \frac{\langle n_i \rangle}{\Omega} = \sum_{r=1}^R S_{ir} \langle f_r(\vec{n}, \Omega) \rangle, \quad (24)$$

$$\partial_t \frac{\langle n_i n_k \rangle}{\Omega^2} = \Omega^{-1} \sum_{r=1}^R S_{kr} \langle n_i f_r(\vec{n}, \Omega) \rangle + S_{ir} \langle n_k f_r(\vec{n}, \Omega) \rangle + S_{kr} S_{ir} \langle f_r(\vec{n}, \Omega) \rangle, \quad (25)$$

$$\begin{aligned} \partial_t \frac{\langle n_i n_k n_l \rangle}{\Omega^3} = & \Omega^{-2} \sum_{r=1}^R [S_{kr} \langle n_i n_l f_r(\vec{n}, \Omega) \rangle + (k \leftrightarrow i) + (i \leftrightarrow l)] \\ & + [S_{ir} S_{kr} \langle n_l f_r(\vec{n}, \Omega) \rangle + (l \leftrightarrow k) + (k \leftrightarrow i)] + S_{kr} S_{ir} S_{lr} \langle f_r(\vec{n}, \Omega) \rangle, \end{aligned} \quad (26)$$

$$\begin{aligned} \partial_t \frac{\langle n_i n_k n_l n_m \rangle}{\Omega^4} = & \Omega^{-3} \sum_{r=1}^R [S_{kr} \langle n_i n_l n_m f_r(\vec{n}, \Omega) \rangle + (k \leftrightarrow i) + (i \leftrightarrow l) + (l \leftrightarrow m)] \\ & + [S_{ir} S_{kr} \langle n_l n_m f_r(\vec{n}, \Omega) \rangle + (i \leftrightarrow m) + (k \leftrightarrow l) + (l \leftrightarrow i)] \\ & + [S_{ir} S_{kr} S_{lr} \langle n_m f_r(\vec{n}, \Omega) \rangle + (m \leftrightarrow i) + (i \leftrightarrow l) + (l \leftrightarrow k)] \\ & + S_{kr} S_{ir} S_{lr} S_{mr} \langle f_r(\vec{n}, \Omega) \rangle, \end{aligned} \quad (27)$$

where  $f_r(\vec{n}, \Omega)$  is the microscopic rate function as defined in Eq. (12). We now assume that the moments can be written as a series in powers of  $\Omega^{-1/2}$ . For the first three moments, for instance, we write

$$\left\langle \frac{n_i}{\Omega} \right\rangle = \sum_{j=0}^4 \alpha_j^i \Omega^{-j/2}, \quad \left\langle \frac{n_i n_k}{\Omega \Omega} \right\rangle = \sum_{j=0}^4 \rho_j^{ik} \Omega^{-j/2}, \quad \left\langle \frac{n_i n_k n_l}{\Omega \Omega \Omega} \right\rangle = \sum_{j=0}^4 \lambda_j^{ikl} \Omega^{-j/2}, \quad (28)$$

where we introduced the coefficients  $\alpha, \rho$  and  $\lambda$ . Note that we have here omitted terms of higher order than  $\Omega^{-2}$ . The subscripts and superscripts indicate expansion orders and species indices, respectively. Substituting Eq. (28) into Eq. (24) and equating equal order terms on both sides of the equation, one obtains:

$$\frac{\partial}{\partial t} \alpha_j^i = \sum_{r=1}^R S_{ir} [\delta_{j,0} \Delta_r + \sum_{k'=1}^N \beta_r^{k'} \alpha_j^{k'} + \sum_{k',s=1, s \neq k'}^N \gamma_r^{k's} \rho_j^{k's} + \sum_{k'=1}^N \zeta_r^{k'k'} (\rho_j^{k'k'} - \alpha_{j-2}^{k'})], \quad j = 0, \dots, 4, \quad (29)$$

where  $\delta_{i,j}$  is the Kronecker delta and  $\alpha_j^{k'} = 0$  if  $j < 0$ . Similarly, one can obtain equations for the coefficients of higher order moments. For instance substituting Eq. (28) into Eq. (25) and equating equal order terms on both sides of the equation, we obtain time-evolution

equations for the coefficients  $\rho_j^{ik}$  of the second order moments:

$$\begin{aligned}
\partial_t \rho_j^{ik} = & \sum_{r=1}^R \left[ S_{kr} \Delta_r(\alpha_j^i) + S_{ir} \Delta_r(\alpha_j^k) \right] + \sum_{r=1}^R \sum_{k'=1}^N \left[ S_{kr} \beta_r^{k'}(\rho_j^{ik'}) + S_{ir} \beta_r^{k'}(\rho_j^{kk'}) \right] \\
& + \sum_{r=1}^R \sum_{k',s=1,s \neq k'}^N \left[ S_{kr} \gamma_r^{k's}(\lambda_j^{ik's}) + S_{ir} \gamma_r^{k's}(\lambda_j^{kk's}) \right] + \sum_{r=1}^R \sum_{k'=1}^N \left[ S_{kr} \zeta_r^{k'k'}(\lambda_j^{ik'k'}) \right. \\
& \left. + S_{ir} \zeta_r^{k'k'}(\lambda_j^{kk'k'}) \right] - \sum_{r=1}^R \sum_{k'=1}^N \left[ S_{kr} \zeta_r^{k'k'}(\rho_{j-2}^{ik'}) + S_{ir} \zeta_r^{k'k'}(\rho_{j-2}^{kk'}) \right] \\
& + \sum_{r=1}^R \sum_{k'=1}^N S_{kr} S_{ir} \left[ \Delta_r \delta_{j,2} + \beta_r^{k'}(\alpha_{j-2}^{k'}) + \sum_{s=1,s \neq k'}^N \gamma_r^{k's}(\rho_{j-2}^{k's}) \right. \\
& \left. + \zeta_r^{k'k'}(\rho_{j-2}^{k'k'} - \alpha_{j-4}^{k'}) \right], \quad j = 0, \dots, 4. \tag{30}
\end{aligned}$$

Note that it is here understood that  $\rho_j^{ks} = 0$  if  $j < 0$ . Equations for the coefficients of third and higher order moments can be similarly derived. As we will show in the following section, Eqs. (29) and Eqs. (30) agree exactly with the ones obtained in the previous section using the conventional SSE.

## V. EQUIVALENCE OF THE TWO METHODS

Comparing Eq. (28) with Eqs. (18)-(20) one obtains the following relationship between the coefficients of the system-size expansion and of the alternative method:

$$\begin{aligned}
\alpha_0^i &= \phi_i, \quad \alpha_{j+1}^i = [\epsilon_i]_j, \quad 0 \leq j \leq 3 \\
\rho_0^{ik} &= \alpha_0^i \alpha_0^k, \quad \rho_1^{ik} = \alpha_1^i \alpha_0^k + \alpha_0^i \alpha_1^k, \quad \rho_{j+2}^{ik} = [\epsilon_i \epsilon_k]_j + \alpha_0^i \alpha_{j+2}^k + \alpha_0^k \alpha_{j+2}^i, \quad 0 \leq j \leq 2 \\
\lambda_0^{ikl} &= \alpha_0^i \rho_0^{kl}, \quad \lambda_1^{ikl} = \alpha_1^i \rho_0^{kl} + (i \leftrightarrow k) + (k \leftrightarrow l), \quad \lambda_2^{ikl} = [\alpha_0^i \rho_2^{kl} - \alpha_2^i \rho_0^{kl}] + (i \leftrightarrow k) + (k \leftrightarrow l), \\
\lambda_{j+3}^{ikl} &= [\epsilon_i \epsilon_k \epsilon_l]_j + [\alpha_0^i \rho_{j+3}^{kl} - \rho_0^{kl} \alpha_{j+3}^i + (i \leftrightarrow k) + (k \leftrightarrow l)], \quad 0 \leq j \leq 1. \tag{31}
\end{aligned}$$

Equivalence of the two methods is then proved if, given these relations, we can obtain the time-dependent equations for the coefficients of the alternative method, i.e., Eqs. (29) and (30), by starting from the time-dependent equations for the coefficients of the SSE, i.e., Eqs. (A1-A10). This is what we seek to establish next.

**Equivalence at the deterministic level.** It follows directly from the previous relations that the deterministic rate equations given by Eq. (15) correspond to Eq. (29) with  $j = 0$ .

**Equivalence at the first moment level.** We start recalling the time evolution equation

for  $[\epsilon_i]_{j'}$  derived using the SSE (see Eqs. (A1)-(A4)):

$$\frac{\partial}{\partial t}[\epsilon_i]_{j'} = J_i^w[\epsilon_w]_{j'} + \left[ \frac{1}{2} J_i^{wp}[\epsilon_w \epsilon_p]_{j'-1} - \frac{1}{2} J_i^{ww}(\phi_w \delta_{j',1} + [\epsilon_w]_{j'-2}(1 - \delta_{j',1})) \right] (1 - \delta_{j',0}), \quad (32)$$

where  $j' = 0, \dots, 3$ . Inserting the relations in Eq. (31) in the above equation, one obtains:

$$\frac{\partial}{\partial t} \alpha_1^i = J_i^w \alpha_1^w, \quad j' = 0, \quad (33)$$

$$\frac{\partial}{\partial t} \alpha_2^i = J_i^w \alpha_2^w + \frac{1}{2} J_i^{wp}(\rho_2^{wp} - (\alpha_0^w \alpha_2^p + \alpha_2^w \alpha_0^p)) - \frac{1}{2} J_i^{ww} \alpha_0^w, \quad j' = 1, \quad (34)$$

$$\frac{\partial}{\partial t} \alpha_3^i = J_i^w \alpha_3^w + \frac{1}{2} J_i^{wp}(\rho_3^{wp} - (\alpha_0^w \alpha_3^p + \alpha_3^w \alpha_0^p)) - \frac{1}{2} J_i^{ww} \alpha_1^w, \quad j' = 2, \quad (35)$$

$$\frac{\partial}{\partial t} \alpha_4^i = J_i^w \alpha_4^w + \frac{1}{2} J_i^{wp}(\rho_4^{wp} - (\alpha_4^w \alpha_0^p + \alpha_4^p \alpha_0^w)) - \frac{1}{2} J_i^{ww} \alpha_2^w, \quad j' = 3. \quad (36)$$

The next step is to substitute the explicit expression of the matrices  $J$  as given by Eq. (23).

This leads to the equations:

$$\begin{aligned} \frac{\partial}{\partial t} \alpha_1^i &= \sum_{r=1}^R \sum_{k'=1}^N S_{ir} \left( \beta_r^{k'} \alpha_1^{k'} + \zeta_r^{k'k'} \rho_1^{k'k'} + \sum_{s=1, s \neq k'}^N \gamma_r^{k's} \rho_1^{k's} \right), \\ \frac{\partial}{\partial t} \alpha_2^i &= \sum_{r=1}^R \sum_{k'=1}^N S_{ir} \left( \beta_r^{k'} \alpha_2^{k'} + \sum_{s=1, s \neq k'}^N \gamma_r^{k's} \rho_2^{k's} + \zeta_r^{k'k'} (\rho_2^{k'k'} - \alpha_0^{k'}) \right), \\ \frac{\partial}{\partial t} \alpha_3^i &= \sum_{r=1}^R \sum_{k'=1}^N S_{ir} \left( \beta_r^{k'} \alpha_3^{k'} + \sum_{s=1, s \neq k'}^N \gamma_r^{k's} \rho_3^{k's} + \zeta_r^{k'k'} (\rho_3^{k'k'} - \alpha_1^{k'}) \right), \\ \frac{\partial}{\partial t} \alpha_4^i &= \sum_{r=1}^R \sum_{k'=1}^N S_{ir} \left( \beta_r^{k'} \alpha_4^{k'} + \sum_{s=1, s \neq k'}^N \gamma_r^{k's} \rho_4^{k's} + \zeta_r^{k'k'} (\rho_4^{k'k'} - \alpha_2^{k'}) \right). \end{aligned} \quad (37)$$

These are exactly the same as Eq. (29) with  $j = 1, \dots, 4$ .

**Equivalence at the second moment level.** Using the SSE we previously derived the expression for  $[\epsilon_i \epsilon_k]_0$  which is given by:

$$\frac{\partial}{\partial t} [\epsilon_i \epsilon_k]_0 = J_i^w [\epsilon_w \epsilon_k]_0 + (i \leftrightarrow k) + J_{ik}. \quad (38)$$

Writing  $[\epsilon_i \epsilon_k]_0$  and  $[\epsilon_w \epsilon_k]_0$  in terms of the new coefficients using Eq. (31), we obtain:

$$\frac{\partial}{\partial t} (\rho_2^{ik} - (\alpha_0^i \alpha_2^k + \alpha_2^i \alpha_0^k)) = [J_i^w (\rho_2^{wk} - (\alpha_0^w \alpha_2^k + \alpha_2^w \alpha_0^k)) + (i \leftrightarrow k)] + J_{ik}. \quad (39)$$

This equation can be expanded as:

$$\frac{\partial}{\partial t} \rho_2^{ik} = \alpha_2^k \frac{\partial}{\partial t} \alpha_0^i + \alpha_0^i \frac{\partial}{\partial t} \alpha_2^k + \alpha_2^i \frac{\partial}{\partial t} \alpha_0^k + \alpha_0^k \frac{\partial}{\partial t} \alpha_2^i + [J_i^w (\rho_2^{wk} - (\alpha_0^w \alpha_2^k + \alpha_2^w \alpha_0^k)) + (i \leftrightarrow k)] + J_{ik}. \quad (40)$$

Next, we substitute the time derivative of  $\alpha_k^i$  (which we derived earlier and is given by Eq. (37)) in the right hand side of the above equation, leading to:

$$\begin{aligned}
\frac{\partial}{\partial t} \rho_2^{ik} = & \alpha_2^k \left( \sum_{r=1}^R \sum_{k'=1}^N S_{ir} \left( \Delta_r + \beta_r^{k'} \alpha_0^{k'} + \sum_{s=1, s \neq k'}^N \gamma_r^{k's} \rho_0^{k's} + \zeta_r^{k'k'} \rho_0^{k'k'} \right) \right) \\
& + \alpha_0^i \left( \sum_{r=1}^R \sum_{k'=1}^N S_{kr} \left( \beta_r^{k'} \alpha_2^{k'} + \sum_{s=1, s \neq k'}^N \gamma_r^{k's} \rho_2^{k's} + \zeta_r^{k'k'} (\rho_2^{k'k'} - \alpha_0^{k'}) \right) \right) \\
& + \alpha_2^i \left( \sum_{r=1}^R \sum_{k'=1}^N S_{kr} \left( \Delta_r + \beta_r^{k'} \alpha_0^{k'} + \sum_{s=1, s \neq k'}^N \gamma_r^{k's} \rho_0^{k's} + \zeta_r^{k'k'} \rho_0^{k'k'} \right) \right) \\
& + \alpha_0^k \left( \sum_{r=1}^R \sum_{k'=1}^N S_{ir} \left( \beta_r^{k'} \alpha_2^{k'} + \sum_{s=1, s \neq k'}^N \gamma_r^{k's} \rho_2^{k's} + \zeta_r^{k'k'} (\rho_2^{k'k'} - \alpha_0^{k'}) \right) \right) \\
& + [J_i^w (\rho_2^{wk} - (\alpha_0^w \alpha_2^k + \alpha_2^w \alpha_0^k)) + (i \leftrightarrow k)] + J_{ik}.
\end{aligned} \tag{41}$$

To write the final explicit equation for  $\rho_2^{ik}$ , one needs to insert the expressions for the  $J$  matrices given by Eqs. (22) - (23). After some algebraic manipulation most of the terms inside the expression cancel leading to the final form:

$$\begin{aligned}
\partial_t \rho_2^{ik} = & \sum_{r=1}^R \left[ S_{kr} \Delta_r (\alpha_2^i) + S_{ir} \Delta_r (\alpha_2^k) \right] + \sum_{r=1}^R \sum_{k'=1}^N \left[ S_{kr} \beta_r^{k'} (\rho_2^{ik'}) + S_{ir} \beta_r^{k'} (\rho_2^{kk'}) \right] \\
& + \sum_{r=1}^R \sum_{k', s=1, s \neq k'}^N \left[ S_{kr} \gamma_r^{k's} (\lambda_2^{ik's}) + S_{ir} \gamma_r^{k's} (\lambda_2^{kk's}) \right] + \sum_{r=1}^R \sum_{k'=1}^N \left[ S_{kr} \zeta_r^{k'k'} (\lambda_2^{ik'k'}) \right. \\
& \left. + S_{ir} \zeta_r^{k'k'} (\lambda_2^{kk'k'}) \right] - \sum_{r=1}^R \sum_{k'=1}^N \left[ S_{kr} \zeta_r^{k'k'} (\rho_0^{ik'}) + S_{ir} \zeta_r^{k'k'} (\rho_0^{kk'}) \right] \\
& + \sum_{r=1}^R \sum_{k'=1}^N S_{kr} S_{ir} \left[ \Delta_r + \beta_r^{k'} (\alpha_0^{k'}) + \sum_{s=1, s \neq k'}^N \gamma_r^{k's} (\rho_0^{k's}) + \zeta_r^{k'k'} (\rho_0^{k'k'}) \right],
\end{aligned} \tag{42}$$

where we used the expressions for the  $\lambda$  coefficients given in Eq. (31). Note that Eq. (42) is precisely one and the same as Eq. (30) with  $j = 2$ .

By a completely analogous derivation to the one above, starting from the SSE time-evolution equations for  $[\epsilon_i \epsilon_k]_1$  and  $[\epsilon_i \epsilon_k]_2$  as given by Eqs. (A6)-(A7), one can derive Eq. (30) with  $j = 3$  and  $j = 4$ , respectively.

Hence in summary we have here proved that the SSE equations for the first two moments of the concentrations up to order  $\Omega^{-2}$  are precisely the same as those obtained from the alternative route. The algebra becomes formidable to perform by hand for third and higher-order moments and for higher powers in  $\Omega$ , but we have carried this out with Mathematica [23] and have verified agreement of the SSE and the alternative method for these cases as well.



We finish this section by noting the difference between our method and moment-closure approximations [12, 24–27]. Essentially the latter close moment equations by postulating an underlying distribution such as Gaussian or log-normal but in contrast in our method no such artificial closure is used. Rather the equations for the moments close by themselves due to the nature of the perturbation expansion. Hence our procedure is unique and does not suffer from the ad-hoc nature of moment-closure approximations. A comparison of the SSE (to which our method is formally equivalent) and of common moment-closure approximations can be found in [28].

## VI. SUMMARY AND CONCLUSION

In this paper we have introduced a novel way to approximately calculate the moments of concentrations in the chemical master equation. This approach is useful when traditional exact methods become computationally demanding, e.g. when one is using the Gillespie algorithm and the number of reactions fired per unit time is large or when one is directly solving the CME equations and the transition matrix has a large dimensionality. Our approach developed in Section IV is based on a Taylor series expansion of the moments of molecule numbers in powers of a small parameter – the inverse square root of the system size. Substituting this expansion in the exact equations for the moments derived from the CME, one obtains time-evolution equations for the coefficients of the Taylor series. We showed in Section V that this approach leads to the same result as the SSE (stated in general in Section III), but in a much simpler and straightforward way. Our approach has one further and most important advantage: unlike the SSE, the moments are not calculated from a GFPE with third and higher-order coefficients and hence it circumvents the problem with the probabilistic interpretation of the GFPE. The fact that it nevertheless agrees with the SSE does prove that the latter is a trustworthy device to approximate the moments.

### Acknowledgements

Claudia Cianci would like to thank Philipp Thomas for useful discussions.

- 
- [1] N. van Kampen, in *Stochastic Processes in Physics and Chemistry (Third Edition)* (Elsevier, Amsterdam, 2007), North-Holland Personal Library.
- [2] D. T. Gillespie, *The journal of physical chemistry* **81**, 2340 (1977).
- [3] L. Cantini, C. Cianci, D. Fanelli, E. Massi, L. Barletti, and M. Asllani, *Journal of mathematical biology* **69**, 1585 (2014).
- [4] J. Elf and M. Ehrenberg, *Genome research* **13**, 2475 (2003).
- [5] P. Thomas, A. V. Straube, and R. Grima, *BMC systems biology* **6**, 39 (2012).
- [6] R. Grima, *Physical Review E* **84**, 056109 (2011).
- [7] R. Grima, P. Thomas, and A. V. Straube, *Journal of Chemical Physics* **8** (2011).
- [8] R. Grima, *The Journal of chemical physics* **133**, 035101 (2010).
- [9] R. Grima, *Physical Review Letters* **102**, 218103 (2009).
- [10] C. Cianci, F. Di Patti, and D. Fanelli, *EPL* (2011).
- [11] C. Cianci, F. Di Patti, D. Fanelli, and L. Barletti, *EPJST* **212**, 5 (2012).
- [12] D. Schnoerr, G. Sanguinetti, and R. Grima, *Journal of Physics A: Mathematical and Theoretical* **50**, 093001 (2017).
- [13] P. Thomas, H. Matuschek, and R. Grima, *PLoS ONE* **6**, e38518 (2012).
- [14] P. Thomas and R. Grima, *Physical Review E* **92**, 012120 (2015).
- [15] R. Ramaswamy, N. Gonzalez-Segredo, I. F. Sbalzarini, and R. Grima, *Nature Communications* **3**, 779 (2012).
- [16] R. F. Pawula, *Physical Review* **162**, 186 (1967).
- [17] H. Risken and H. D. Vollmer., *Z. Physik B* **35**, 313 (1979).
- [18] W. Horsthemke and L. Brenig, *Z. Physik B* **27**, 341 (1977).
- [19] D. R. Glowacki, C.-H. Liang, C. Morley, M. J. Pilling, and S. H. Robertson, *Journal of Physical Chemistry A* **116**, 9545 (2012).
- [20] A. J. McKane and T. J. Newman, *Physical Review Letters* **94**, 218102 (2005).
- [21] P. Thomas, Ph.D. thesis, University of Edinburgh (2015).
- [22] C. M. Bender and S. A. Orszag, *Advanced Mathematical Methods for Scientist and Engineers* (Springer, 1999).
- [23] *Mathematica*, (Wolfram Research, Inc.) (2015).

- [24] A. Singh and J. Hespanha, 45th IEEE Conference on Decision and Control (2006).
- [25] A. Ale, P. Kirk, and M. Stumpf, The Journal of chemical physics **17**, 174101 (2013).
- [26] D. Schnoerr, G. Sanguinetti, and R. Grima, The Journal of chemical physics **141**, 024103 (2014).
- [27] D. Schnoerr, G. Sanguinetti, and R. Grima, The Journal of chemical physics **18**, 185101 (2015).
- [28] R. Grima, Journal of Chemical Physics **136**, 154105 (2012).

### Appendix A: Time-evolution equations for the pseudo-moments of the SSE

For the computation of the average concentrations up to order  $\Omega^{-2}$  we need  $[\epsilon_i]_j$ ,  $j = 0, \dots, 3$ , (see Eq. (18)) which are given by the equations:

$$\frac{\partial}{\partial t} [\epsilon_i]_0 = J_i^w [\epsilon_w]_0, \quad (\text{A1})$$

$$\frac{\partial}{\partial t} [\epsilon_i]_1 = J_i^w [\epsilon_w]_1 + \frac{1}{2} J_i^{wp} [\epsilon_w \epsilon_p]_0 - \frac{1}{2} J_i^{ww} \phi_w, \quad (\text{A2})$$

$$\frac{\partial}{\partial t} [\epsilon_i]_2 = J_i^w [\epsilon_w]_2 + \frac{1}{2} J_i^{wp} [\epsilon_w \epsilon_p]_1 - \frac{1}{2} J_i^{ww} [\epsilon_w]_0, \quad (\text{A3})$$

$$\frac{\partial}{\partial t} [\epsilon_i]_3 = J_i^w [\epsilon_w]_3 + \frac{1}{2} J_i^{wp} [\epsilon_w \epsilon_p]_2 - \frac{1}{2} J_i^{ww} [\epsilon_w]_1. \quad (\text{A4})$$

According to Eq. (19), for the computation of the second moments of the concentrations up to order  $\Omega^{-2}$ , we need  $[\epsilon_i]_j$ ,  $j = 0, \dots, 3$ , and  $[\epsilon_i \epsilon_l]_j$ ,  $j = 0, \dots, 2$ . The latter are given by the equations:

$$\frac{\partial}{\partial t} [\epsilon_i \epsilon_k]_0 = J_i^w [\epsilon_w \epsilon_k]_0 + (i \leftrightarrow k) + J_{ik}, \quad (\text{A5})$$

$$\frac{\partial}{\partial t} [\epsilon_i \epsilon_k]_1 = J_i^w [\epsilon_w \epsilon_k]_1 - \frac{1}{2} J_i^{ww} [\epsilon_k]_0 \phi_w + \frac{1}{2} J_i^{wp} [\epsilon_w \epsilon_p \epsilon_k]_0 + (i \leftrightarrow k) + J_{ki}^w [\epsilon_w]_0, \quad (\text{A6})$$

$$\begin{aligned} \frac{\partial}{\partial t} [\epsilon_i \epsilon_k]_2 &= J_i^w [\epsilon_w \epsilon_k]_2 + \frac{1}{2} J_i^{wp} [\epsilon_w \epsilon_p \epsilon_k]_1 - \frac{1}{2} J_i^{ww} \phi_w [\epsilon_k]_1 \\ &\quad - \frac{1}{2} J_i^{ww} [\epsilon_w \epsilon_k]_0 + (i \leftrightarrow k) + J_{ik}^w [\epsilon_w]_1 + \frac{1}{2} J_{ik}^{wm} [\epsilon_w \epsilon_m]_0 - \frac{1}{2} J_{ik}^{ww} \phi_w. \end{aligned} \quad (\text{A7})$$

We observe that the equations for  $[\epsilon_w \epsilon_k]_1$  and  $[\epsilon_w \epsilon_k]_2$  depend on the zeroth and first order coefficients  $[\epsilon_w \epsilon_p \epsilon_k]_0$  and  $[\epsilon_w \epsilon_p \epsilon_k]_1$  of the third moment whose time-evolution equations are

given by:

$$\frac{\partial}{\partial t}[\epsilon_i \epsilon_k \epsilon_l]_0 = J_l^w[\epsilon_w \epsilon_i \epsilon_k]_0 + (l \leftrightarrow k) + (k \leftrightarrow i) + J_{il}[\epsilon_k]_0 + (k \leftrightarrow i) + (i \leftrightarrow l), \quad (\text{A8})$$

$$\begin{aligned} \frac{\partial}{\partial t}[\epsilon_i \epsilon_k \epsilon_l]_1 &= J_l^w[\epsilon_w \epsilon_k \epsilon_i]_1 + \frac{1}{2} J_l^{wp}[\epsilon_w \epsilon_p \epsilon_i \epsilon_k]_0 - \frac{1}{2} J_l^{ww} \phi_w[\epsilon_i \epsilon_k]_0 + (l \leftrightarrow i) + (i \leftrightarrow k) \\ &+ J_{il}[\epsilon_k]_1 + J_{il}^w[\epsilon_w \epsilon_k]_0 + (k \leftrightarrow i) + (i \leftrightarrow l) + J_{ikl}. \end{aligned} \quad (\text{A9})$$

Note that according to our previously introduced shorthand notation, the terms  $J_{il}[\epsilon_k]_0 + (k \leftrightarrow i) + (i \leftrightarrow l)$  in Eq. (A8) stand for  $J_{il}[\epsilon_k]_0 + J_{kl}[\epsilon_i]_0 + J_{ki}[\epsilon_l]_0$ .

Finally we note that the last equation involves the zeroth order coefficient of the fourth moment, which satisfies the time-evolution equation:

$$\begin{aligned} \frac{\partial}{\partial t}[\epsilon_i \epsilon_k \epsilon_l \epsilon_m]_0 &= J_i^w[\epsilon_w \epsilon_k \epsilon_l \epsilon_m]_0 + (i \leftrightarrow m) + (m \leftrightarrow k) + (k \leftrightarrow l) + J_{im}[\epsilon_k \epsilon_l]_0 + (m \leftrightarrow l) \\ &+ (l \leftrightarrow k) + (i \leftrightarrow m) + (m \leftrightarrow l) + (k \leftrightarrow m). \end{aligned} \quad (\text{A10})$$

## Appendix B: List of acronyms

SSA stochastic simulation algorithm

SSE system-size expansion

LNA linear-noise approximation

GFPE generalised Fokker-Planck equation

CME chemical master equation